

This manuscript has been accepted after a double-blind peer review and will be published in a future issue of this journal. Its content is final and only the information regarding volume, number, publication date and pagination will change in the final version

Potential uses of *Ustilago maydis*

Potenciales usos de *Ustilago maydis*

Isi I. Ordóñez Velázquez^a

Abstract:

Due to the high biodiversity of ecosystems in Mexico, there is a high variety of species, some of the most abundant are fungi, since there is an approximate record of 200 thousand endemic species of Mexico, traditionally the use given to these organisms is food and medicinal. One of the most characteristic species of our country is *Ustilago maydis* commonly known as huitlacoche, which has various properties in its high nutritional value in amino acids, fats, antioxidants, proteins and its ancestral use to treat different painful conditions. These applications have fostered the interest of different industries such as pharmaceuticals, since different studies show that *U. maydis* could have potential biotechnological uses. For this reason, this narrative review article aims to disseminate relevant biological aspects and the potential uses of huitlacoche.

Keywords:

Huitlacoche, medicinal mushrooms, antioxidant, edible mushrooms, nutrients

Resumen:

Debido a la alta biodiversidad de ecosistemas en México, existe una alta variedad de especies, unas de las más abundantes son hongos, ya que se tiene un registro aproximado de 200 mil especies endémicas en México, tradicionalmente el aprovechamiento que se le da a estos organismos es alimenticio y medicinal. Una de las especies más características de nuestro país es *Ustilago maydis* comúnmente conocido como huitlacoche, el cual presenta diversas propiedades en su alto valor nutricional en aminoácidos, grasas, antioxidantes, proteínas y su uso ancestral para tratar diferentes padecimientos dolorosos. Estas aplicaciones han fomentado el interés de diferentes industrias como la farmacéutica, ya que diferentes estudios demuestran que *U. maydis* podría tener potenciales usos biotecnológicos. Por tal motivo, este artículo de revisión narrativa pretende difundir aspectos biológicos relevantes y los potenciales usos del huitlacoche.

Palabras Clave:

Huitlacoche, hongos medicinales, antioxidantes, hongos comestibles, nutrientes

INTRODUCTION

Mexico is classified as a diverse country, occupying the fifth place in the world due to its wide number of species and endemisms, it is estimated that it represents 10% of the terrestrial diversity of the planet, due to its geographical location, topography, diversity of altitudes and climates, resulting in the diversification of habitats and different forms of life.¹⁻³ A clear example of this is fungi, which are considered to be the second most numerous organisms on Earth after insects⁴, Mexico has temperate forests of gymnosperms and angiosperms, favoring the development of approximately 200,000 species of fungi of which only 3.5% to 5% have been reported including national and international research⁴⁻⁶, among

the uses and customs are gastronomic and medicinal. In pre-Hispanic times these were used to cure diseases of a spiritual nature such as the evil eye and witchcraft practices (*Dictyophora* and the lycoperdaceae)⁷, currently, there has been an increased interest in modern pharmacology in the extraction and studies of chemical compounds from medicinal edible mushrooms.^{8,9} Among the properties attributed to them are anti-inflammatory, antioxidant, treating different types of pain such as headache, low back pain, among others.¹⁰

The endemic species of Mexico *Ustilago maydis*, is a basidiomycete parasitic fungus known in the world to cause the disease called common corn smut (*Zea mays*), this infection generates large tumors in the cob, culturally known as

^a Consultorio privado | Pachuca | Hidalgo | México, <https://orcid.org/0009-0007-4051-9156>, Email: isiorvel@gmail.com

“huitlacoche” or “cuitlacoche” this fungus is used in the preparation of different dishes¹¹, however, the interest in this species also lies in the potential uses for its biochemical properties.

HOW IS HUITLACOCHÉ PRODUCED?

Fungi are eukaryotic organisms, heterotrophs, chemoorganotrophs since they need organic compounds as the main source of energy, the vast majority of fungi take their nutrients by absorption since they release degrading exoenzymes of complex compounds, which allows them to use a great diversity of substrates as nutrients and energy sources. Filamentous fungi have a basic structure called a hyphae, it is a tubular structure that branches as the fungus grows in an apical form, the set of hyphae that produces the fungus is called micelio.¹²⁻¹⁴ *U. maydis* is a basidiomycete fungus, phytopathogenic, this species is dimorphic with a non-pathogenic yeast form and a pathogenic filamentous form which is responsible for the formation of tumors in corn, which makes it a non-obligated parasite, due to the above this species depends on a living tissue for its proliferation and development^{15,16}, thus the symbiotic relationship between these two species (*U. maydis* and *Zea mays*) is merely parasitic since in this correlation only one species is benefited.¹⁷ *U. maydis* requires one of its two hosts to complete its sexual cycle.¹⁸⁻²⁰ During its life cycle it can be divided into two phases, one of them saprophytic during which the fungus grows in the form of yeasts which, they reproduce by spores, the second phase is mycelial and pathogenic, where the fungus grows in the form of hyphae which only develop inside the host, most of the growth of the fungus occurs in meristematic tissues (embryonic tissue, poorly differentiated cells, responsible for growth and regeneration) of the plant²¹, resulting in galls or tumors where the sporulation of the fungus subsequently occurs. The hyphae grow irregularly and fragment, covered with a thick, equinulate and pigmented wall. Symptoms in the plant can range from tumor formation, chlorosis, distortions to plant dwarfism.^{18,22} The production of huitlacoche can occur in two ways, natural and inoculated. The natural way is the “contamination of the cob” with the fungus strain without human intervention (Figure 1), the artificial way is the inoculation of each of the ears by means of an injection with the strain, carried out by farmers¹¹, it is required that the maize variety (host) be susceptible to the fungus, that the pathogen strain is virulent and the environment is favourable, with 80–85% relative humidity and a temperature of 16–32°C.²³

The huitlacoche in Mexico is not considered a pest, this species can naturally infect 1 to 5% of the crop, however, this percentage gives an economic gain to the farmer since the infected cob is marketable and many times its value is usually higher than that of the unparasitized cob.²²

BIOACTIVE COMPOUNDS

Huitlacoche has been studied as a biofunctional food, which is composed of multiple vitamins (C and E), minerals that are

biologically important for humans²⁴, phenolic compounds and flavonoids which exert antioxidant activity, as they protect from the action of cellular oxidizing species²⁵, the consumption of these compounds are associated with a lower incidence of chronic diseases such as diabetes mellitus and cardiovascular diseases.²⁶ The fatty acids (FAs) present in this species such as oleic and linoleic acids, are precursors of Omega 3 and Omega 6, which are essential for humans, since their deficiency can produce serious metabolic and structural alterations at the cellular level.²⁷




Figure 1. Life cycle without human intervention of *Ustilago maydis*.¹⁸⁻²⁰

These FAs provide fluidity to cell membranes, which is essential for membrane proteins (ion channels, receptors, communicating junctions, catalytic receptors, enzymes, vesicle-forming structures, etc.) to have the mobility they require to perform their functions. It has been shown that in the formation of nervous tissue, particularly the brain, the fluidity of the membranes is especially important.²⁷ The high content of essential fatty acids suggests a high nutritional value of huitlacoche, could be due to the fact that corn is one of the cereals with the highest content of fats and essential unsaturated fatty acids.^{27,28}

The most commonly reported minerals are phosphorus, magnesium, zinc, iron, and sodium being metabolically indispensable for the body^{25,28}, it contains almost all the essential amino acids such as: lysine (most abundant amino acid), glycine, leucine, glutamic acid, aspartic acid, valine, isoleucine, phenylalanine, alanine, serine, tyrosine, proline, threonine, methionine, ornithine, tryptophan, γ -aminobutyric acid, ornithine and trichotomic acid.^{25,29-31}

Regarding the protein value, it has been shown to have a considerable amount of protein in conjunction with other edible mushroom species (Table 1)³²⁻³⁷, *Ustilago maydis* increases the

percentage of protein in corn.^{25,37} This mushroom provides advantages for the consumer over animal proteins by presenting antioxidant properties, antitumor, and antimicrobials. In the industry, protein concentrates, hydrolysates and fungal peptides are made to contribute to improving human health, in addition, edible mushrooms can be used to enrich traditional foods by increasing the protein value and functional qualities of the food, another important factor is the low-fat content, high fiber content³⁸, so it could be considered an alternative and complement in vegetarian diets.²⁸

Table 1 Protein content of edible mushrooms

Species	Protein amount*	References
<i>Agaricus bisporus</i>	19%	32,33
<i>Boletus edulis</i>	18.5	34
<i>Pleurotus ostreatus</i>	17.5%	35,37
<i>Ustilgo maydis</i>	13.46%	36
<i>Amanita caesarea</i>	13%	34

*Crude protein in dry matter

CONSUMPTION OF HUITLACOCHÉ FOR INFLAMMATION AND PAIN

The International Association for the Study of Pain (IASP) defines pain as an unpleasant sensory and/or emotional experience associated with actual or potential tissue damage^{39,40}, which can be classified depending on its duration, pathogenesis, location, course and intensity⁴¹, as a result of cell damage, a process of inflammation is generated, which is a physiological response in order to repair damaged tissue; the inflammatory process begins when chemical compounds are released by the damaged tissue, in response, white blood cells produce substances that cause cells to divide and grow to rebuild tissue and thus repair the injury.

There are two types of inflammation, acute inflammation, which occurs in a short period of time and symptoms last for days, and chronic inflammation, which is described as slow and prolonged, which can last from months to years.⁴² Inflammation can be due to different causes which can range from a blood clot inducing a stroke, immune system disorder, cancer, chemical exposure to polycyclic aromatic hydrocarbons, dioxins, smoking, physical injury including trauma, or a neurological condition.^{43,44} Conditions of inflammation and pain can become disabling for those who suffer from it, impairing quality of life⁴⁵, however, it has been shown that prolonged use of Nonsteroidal anti-inflammatory drugs (NSAIDs) used to treat pain, fever and other inflammatory processes⁴⁶, such as naproxen, ibuprofen, ketorolac, can lead to adverse effects on the cardiovascular, gastrointestinal, hepatic and renal systems.⁴⁷ Due to the above, there has been increased interest in offering alternatives or adjuvants to treat pain, demonstrating the efficacy and biosafety of the use of fungi and medicinal plants through research in biochemical and pharmacological analyses.⁴⁸ Generationally, *Ustilago maydis* has been used in

painful and inflammatory situations such as intestinal pain, muscle inflammation and headaches^{33,49,50}, an ancestral use it is to make a poultice, which is placed on a cloth, and then applied directly to the area of inflammation⁵¹, due to this, interest has been aroused in studying the compounds responsible for mitigating these discomforts; of the compounds recently evaluated for inflammatory and painful processes are ustilagols.³⁰ Ustilagols are compounds extracted from the mycelium of *U. maydis* that exhibit dopaminergic properties, offering pharmacological potential against some neuroleptic diseases⁵⁰, on the other hand, Ustilago C, E demonstrated anti-inflammatory properties, Ustilago G showed that it can generate antiplatelet, similar to that observed with Aspirin®, Ustilagoes A – F demonstrated *in vitro* studies its potent anti-inflammatory and antithrombotic properties.⁵¹ However, one cannot dismiss the ancestral knowledge in which it has been described that huitlacoche has been used to cure heart disease, colic, blisters, pimples, skin burns, athlete's foot, wounds, nosebleeds, rashes in babies, stop bleeding, heal animal bites, relieve dehydration and help with anxiety.^{7,52-55} Nonetheless, there are still areas of opportunity to verify its multiple applications in pain and Inflammation and a possible future as an adjunct to a pharmacological treatment.

ANTIOXIDANT PROPERTIES

The antioxidant capacity of foods is derived from the cumulative synergistic action of a wide variety of antioxidants, such as polyphenols, flavonoids, vitamins C and E, carotenoids, terpenoids, Maillard compounds, and trace elements. These antioxidants play a role in the prevention of diseases related to oxidative⁵⁶⁻⁵⁸, such as neurodegenerative diseases, in which progressive degeneration and neuronal death take place⁵⁹, due to an imbalance in the generation of oxidizing species of nucleic acids, proteins, and lipids resulting in mitochondrial dysfunction, glial cell activation, apoptosis, protein oligomerization, inflammatory response, alteration of the blood-brain barrier⁶⁰⁻⁶², however, phenolic compounds, flavonoids, and carotenoids have been shown to be antioxidant compounds that have potential health benefits, a potential industrial, pharmaceutical, and food application.²⁵

According to National Institute of Statistics and Geography (INEGI) and National Population Council (CONAPO) the trend of the Mexican adult population is increasing⁶³, due to the increase in people's longevity, it is expected that there will be an increase in conical degenerative diseases⁶⁴, adding to this an unbalanced diet with an increase in the consumption of highly processed foods reducing traditional ethnic foods with high nutritional value.⁶⁵

The consumption of antioxidants may be affected; therefore, it is opportune to encourage the Mexican population to consume foods with content of antioxidants. One of these foods is Huitlacoche (table 2)⁶⁶⁻⁷⁰, this food when heat-treated (cooking as food) increases the concentration of phenolic compounds and flavonoids, benefiting its antioxidant capacity, therefore this

food can contribute to an improvement in the state of human health, helping to reduce the damage generated during the oxidative stress of some metabolic processes.^{24,55}

HUITLACOCHES AS FOOD

With the information that is available from prehistory to the present times, the huitlacoche is used as a food in Mexico, mainly in the center and southeast, because of this there is a great variety of ways of preparing this fungus, the National Autonomous University of Mexico published a book entitled 'Cujtlacoche el huitlacoche', an open-access version describes What is it?, How is it made?, trade, life cycle, traditions, history and phylogeny of this species; this book includes approximately 31 recipes ranging from Avocados stuffed with Cuitlacoche (Figure 2) to Salmon with sweet cuitlacoche sauce,⁷¹ in this way, the versatility of this species as a food is shown. With all the scientific support that has been given to huitlacoche, it is considered an excellent food, the best way to take advantage of its properties is cooked^{24,55}, it is important to emphasize that the low use of oil is essential in any dish, in this way it is considered a complement in food plans and highly adaptable to different lifestyles, so promoting its consumption is important.

Table 2 Antioxidant compounds recorded in huitlacoche.

Group	Derived compound	Reference
Phenolics	ferulic	24,28,55,66,67
	Sinapic	
	Chlorogenic	
	p-coumaric	
Flavonoids	caffeic	25,28,55,67
	Anthocyanins	
	Quercetin	
	Naringenin	
	Catechin	
Carotenoids	β -carotene	67,68
	β -cryptoxanthin	
Phytosterols	campesterol-3- β -glucoside	28,69,70
	Δ 7-stigmasterol	
	Δ 7-avenasterol	
	Ergosterol	

FUTURE OF MUSHROOMS

As mentioned above, most mushrooms have a broad profile of essential amino acids, which can contribute to different dietary requirements, while fungi have economic advantages over plant and animal sources.⁷² With the passage of time the food industry seeks to meet the demand for food. It is predicted that by 2050 the world's population will exceed 9 billion people. Agriculture Organization (FAO) says that agricultural production will have to increase food production by 70% to meet demand.⁷³ Due to the nature of fungi, they can be produced industrially in a high quantity and in a short period of time. Within the biotechnological field, multiple species of fungi have been used for processes such as bioconversion, biorefining, bioremediation, biodegradation, antimicrobial and antifungal

agents, although *Ustilago maydis* is more studied to be a complementary and functional food for the population, its biotechnological potential cannot be discarded, since it was shown that it can be used as a complement in biological control for wine production at the beginning of fermentation, since *U. maydis* produces an effective toxin against *Brettanomyces bruxellensis* yeast deteriorating.⁷⁴



Figure 2. A, B: Raw Huitlacoche for sale at the Pachuca Hidalgo supply center, C: Avocados stuffed with huitlacoche and cheese (Recipe based on the book Cujtlacoche el huitlacoche).⁷¹

The antimicrobial and antifungal activity of this species is relevant since its extracts have been shown to be effective against yeasts, bacteria and molds, its glycolipids showed efficacy against *Staphylococcus aureus* and *Salmonella enterica*, *Typhimurium*, *Aspergillus terreus* and *Candida albicans*, which shows that *U. maydis* has potential application in the food, pharmaceutical and industrial industries.^{30,75-77} As has been shown in different molecular studies, fungi are strictly evolutionarily related to animal cells, consequently many of the cellular processes are conserved between these groups; cell model systems of fungi such as yeasts have contributed to the understanding of different cellular pathways, as they have a good genetic accessibility, a short generation time and simple culture form, the role that species like *U. maydis* is that they have an advantage as a yeast. *U. maydis* was shown to have an unexpected similarity in the engines of microtubule organization, long-distance transport and mitosis, with higher eukaryotic cells, this discovery demonstrates that *U. maydis* has

a large number of genes closely related to human genes, this supports the use of this species as a model system for the study of cellular processes that may favor the genetic study of our species.⁷⁸

CONCLUSION

As has been demonstrated through the generations, huitlacoche has a wide variety of uses, from gastronomic to medicinal. It is important to bear in mind that there are still areas of opportunity in which different investigations can be carried out to demonstrate and rectify such uses. The interest in this species in different industries such as pharmaceutical, food and biotechnology have shown that this species is being considered as a good model to study different cellular metabolic processes and at the same time has compounds that could have pharmaceutical potential, although there is still much to be investigated, however, with the already existing evidence. The huitlacoche is undoubtedly a good complementary food for the Mexican diet, taking into account its multiple benefits due to antioxidants, fats and proteins that it contains, adding that the preparation of this is a representative dish in Mexican culture.

REFERENCES

- [1] Koleff P, Urquiza-Haas T. Planeación para la conservación de la biodiversidad terrestre en México: retos en un país megadiverso. México: CONABIO, CONANP; 2011.
- [2] CONABIO. Expertos evalúan y enlistan hongos en riesgo de extinción de los bosques templados de México. 2019.
- [3] Plascencia RL, Castañón Barrientos A, Raz-Guzmán A. La biodiversidad en México su conservación y las colecciones biológicas. Ciencias 2011; 101: 36-43.
- [4] López Sánchez CO. Tierra de Hongos. México: CONABIO; 2022.
- [5] Aguirre-Acosta E, Ulloa M, Aguilar S, Cifuentes J, Valenzuela R. Biodiversidad de hongos en México. Rev. Mex. Biodiv. 2014; 85(Suppl): 76-81.
- [6] Sánchez JE, Mata G. Hongos comestibles y medicinales en Iberoamérica investigación y desarrollo en un entorno multicultural. ed 1. Chiapas México: El colegio de la frontera Sur. 2012.
- [7] Guzmán G. Diversity and use of traditional Mexican medicinal fungi A Review. Int J. Med. Mushrooms 2008; 10(3): 209-17.
- [8] Barros L, Cruz T, Baptista P, Estevinho LM, Ferreira IC. Wild and commercial mushrooms as source of nutrients and nutraceuticals. Food Chem. Toxicol. 2008; 46(8):27 42-7.
- [9] Yu-Cheng D, Zhu-Liang Y, Bao-Kai C, Chang-Jun Y, Li-Wei Z. Species diversity and utilization of medicinal mushrooms and fungi in China (Review). Int. J. Med. Mushrooms 2009; 11(3): 287-302.
- [10] Jiménez Ruiz M, Pérez-Moreno J, Juan J, Almaraz-Suárez, Torres-Aquino M. Hongos silvestres con potencial nutricional, medicinal y biotecnológico comercializados en Valles Centrales, Oaxaca. Rev. Mex. Cienc. Agríc. 2013; 4(2): 199-213.
- [11] Aguayo-González DJ, Guevara-Lara F, Luna-Ruiz JJ, Pérez-Cabrera LE, García Munguía-CA, García-Munguía AM. Patogenicidad de cepas de Ustilago maydis para la producción en condiciones controladas. Rev. Mex. Cienc. Agríc. 2021; 12(3): 513-24.
- [12] Estrada Salazar GI, Ramírez Gelando MC. Micología general. Manizales Colombia: Centro Editorial Universidad Católica de Manizales; 2019.
- [13] Suárez-Contreras LY, Peñaranda-Figueroa FA. Identificación molecular de hongos filamentosos y su potencial biotecnológico. Rev. Bio. Agro. 2022; 20(1): 194-206.
- [14] Cepero de García MC, Restrepo Restrepo S, Franco Molano AE, Cárdenas Toquica M, Vargas Estupiñán N. Biología de hongos. Bogotá Colombia: Ediciones Uniandes; 2012.
- [15] Macuil-Tlachino V, Sobal-Cruz M, Martínez-Carrera D, Morales-Almora P, Peña-Olvera B, Maimone-Celorio MR. Obtención de cepas infectivas de Ustilago maydis para la producción de huitlacoche en la sociedad rural mexicana. Agri. Soc. Desarrollo 2021; 18(3): 335-45.
- [16] Yu C, Qi J, Han H, Wang P, Liu C. Progress in pathogenesis research of Ustilago maydis, and the metabolites involved along with their biosynthesis. Mol. Plant Pathol. 2023; 24(5): 495-509.
- [17] Brefort T, Doehlemann G, Mendoza-Mendoza A, Reissmann S, Djamei A, Kahmann R. Ustilago maydis as a Pathogen. Annu. Rev. Phytopathol. 2009; 47: 423-45.
- [18] Banuett F, Herskowitz I. Discrete developmental stages during teliospore formation in the corn smut fungus, Ustilago maydis. Development 1996; 122(10): 2965-76.
- [19] León-Ramírez CG, Sánchez-Arreguín JA, Ruiz-Herrera J. Ustilago maydis, a Delicacy of the Aztec Cuisine and a Model for Research. Natural Resources 2014; 5: 256-67.
- [20] Ruiz-Herrera J, León-Ramírez C, Martínez-Espinoza AD. Morphogenesis and pathogenesis in Ustilago maydis. Develop. Microbiol. 2000; 5: 585-98.
- [21] Cordero del Campillo M, Rojo Vázquez FA, Martínez Fernández AR: Sánchez Acedo MC, Hernández Rodríguez S, Navarrete López-Cozar I, et al. Parasitología veterinaria: Madrid España; McGraw-Hill Interamericana; 2007.
- [22] Ruiz J, Ustilago maydis: ascenso de un hongo mexicano de la gastronomía local al mundo científico. Nova Scientia 2008; 1(1): 118-35.
- [23] Martínez L, Villanueva C, Sahagún J. Susceptibility and resistance of maize to the edible fungus huitlacoche (Ustilago maydis Cda.) improving its virulence. Rev. Chapingo Serie Hort. 2000; 6 (2): 241-48.
- [24] López-Martínez LX, Aguirre-Delgado A, Saenz-Hidalgo HK, Buenrostro-Figueroa JJ, García HS, Baeza-Jiménez R. Bioactive ingredients of huitlacoche (Ustilago maydis), a potential food raw material. Food Chem. (Oxf) 2022; 22(4): 100076.
- [25] Valdez-Morales M, Barry K, Fahey JrGC, Domínguez J, Gonzalez de Mejia E, Valverde ME. Effect of maize genotype, developmental stage, and cooking process on the nutraceutical potential of huitlacoche (Ustilago maydis). Food Chem. 2010; 119(2): 689-97.
- [26] Cereceres-Aragón A, Rodrigo-García J, Álvarez-Parrilla E, Rodríguez-Tadeo A. Ingestión de compuestos fenólicos en población adulta mayor. Nutr. Hosp. 2019; 36(2): 470-8.
- [27] Molina Montes E, Martín Islán AP. Ácidos grasos esenciales Omega-3 y Omega-6 papel en el embarazo y la lactancia. Of. 2010; 29(1): 66-72.
- [28] Aydoğdu M, Gölükçü M. Nutritional value of huitlacoche, maize mushroom caused by Ustilago maydis. Food Sci. Technol. 2017; 37(4): 531-5.

- [29] Méndez-López A, Sánchez-Vega M, Cruz-Salazar Torres J, Martínez-Amador YS, Leal-Robles A. El cultivo de huitlacoche: alimento nutritivo y sustentable. *Rev. Cien. Agro. Apli. Biotec.* 2019; 9-12.
- [30] Villagrán Z, Martínez-Reyes M, Gómez-Rodríguez H, Ríos-García U, Montalvo-González E, Ortiz-Basurto RI, et al. Huitlacoche (*Ustilago maydis*), an iconic Mexican fungal resource: biocultural importance, nutritional content, bioactive compounds and potential biotechnological applications. *Molecules* 2023; 29;28(11): 4415.
- [31] Martínez-Flores A, Corrales-García JJ, Espinosa-Solares T, García-Gatica PG, Villanueva-Verduzco C. Postharvest changes in the edible mushroom huitlacoche (*Ustilago maydis* (D.C.) Corda). *Rev. Chapingo Ser. Hortic.* 2008; 14(3): 339-46.
- [32] Bernás E, Jaworska G, Lisiewska Z. Edible mushroom as a resource of valuable nutritive constituents. *Acta Sci. Pol. Technol. Aliment.* 2006; 5(1): 5-20.
- [33] Cano-Estrada A, Romero-Bautista L. Valor económico, nutricional y medicinal de hongos comestibles silvestres. *Rev. Chil. Nutr.* 2016; 43(1): 75-80.
- [34] Instituto Politecnico Nacional. Catálogo de hongos comestibles de Santiago Papasquiaro. Sistema de Administración de Programas y Proyectos de Investigación. IPN 2002; 1-27.
- [35] Bonatti M, Karnopp P, Soares HM, Furlan SA. Evaluation of *Pleurotus ostreatus* y *Pleurotus sajor-caju* nutricional characteristics when cultivated in different lignocellulosic waste. *Food Chemistry* 2004; 88(3): 425-28.
- [36] Pimentel-González DJ, Rodríguez-Huezo ME, Campos-Montiel RG, Trapala-Islas A, Hernández-Fuentes AD. Influencia de la variedad de maíz en las características fisicoquímicas del huitlacoche (*Ustilago maydis*). *Rev. Mex. Ing. Quim.* 2011; 10(2): 171-8.
- [37] Vanegas PE, Valverde ME, Paredes-López O, Pataky JK. Production of the Edible Fungus Huitlacoche (*Ustilago maydis*): Effect of Maize Genotype on Chemical Composition. *J. Ferm. Bioen.* 1995; 80(1): 104-6
- [38] Ayimbila F, Keawsompong S. Nutritional quality and biological application of mushroom protein as a novel protein alternative. *Curr. Nutr. Rep.* 2023; 12(2): 290-307.
- [39] Krenzischek DA, Wilson L, Newhouse R, Mamaril M, Kane HL. Clinical evaluation of the ASPAN pain and comfort clinical guideline. *J. Perianesth. Nurs.* 2004; 19(3): 150-63.
- [40] Treede RD, Rief W, Barke A, Aziz Q, Bennett MI, Benoliel R, et al. Chronic pain as a symptom or a disease: the IASP classification of chronic pain for the international classification of diseases (ICD-11) *Pain* 2019; 160(1): 19-27.
- [41] Anekar AA, Maxwell Hendrix JM, Cascella M. WHO Analgesic Ladder. Treasure Island: StatPearls; 2024.
- [42] Pahwa R, Goyal A, Jialal I. Chronic inflammation. Treasure Island: StatPearls; 2023.
- [43] Doll DN, Barr TL, Simpkins JW. Cytokines: their role in stroke and potential use as biomarkers and therapeutic targets. *Aging Dis.* 2014; 5(5): 294-306.
- [44] Banerjee A, Khemka VK, Roy D, Dhar A, Sinha Roy TK, Biswas A, et al. Role of pro-inflammatory cytokines and vitamin D in probable Alzheimer's disease with depression. *Aging Dis.* 2017; 8(3): 267-6.
- [45] Woolf CJ. Central sensitization: implications for the diagnosis and treatment of pain. *Pain* 2011; 152(3 Suppl): S2-S15.
- [46] Ghlichloo I, Gerriets V. Nonsteroidal Anti-inflammatory drugs (NSAIDs). Treasure Island: StatPearls; 2024.
- [47] Dávila Cabo de Villa E, Morejón Hernández JM, Acosta Figueredo E. Pain and painkillers. some timely considerations. *Medisur* 2020; 18(4): 694-705.
- [48] Quiñonez-Bastidas GN, Navarrete A. Mexican plants and derivatives compounds as alternative for inflammatory and neuropathic pain treatment-A Review. *Plants (Basel)* 2021; 10(5): 865.
- [49] Venturella G, Ferraro V, Cirlincione F, Gargano ML. Medicinal mushrooms: Bioactive compounds, use, and clinical trials. *Int. J. Mol. Sci.* 2021; 22(2): 634.
- [50] Kurz M, Eder C, Isert D, Li Z, Paulus EF, Schiell M, et al. Ustilipids, acylated β -D-mannopyranosyl D-erythritols from *Ustilago maydis* and *Geotrichum candidum*. *J. Antibiot. (Tokio)* 2003; 56(2): 91-101.
- [51] Hope ME, Pereyra L. Nuestro maíz, treinta monografías populares. México: Museo nacional de culturas populares; 1982.
- [52] Wu HC, His HY, Hsiao G, Yen CH, Leu JY, Wu CC et al. Chemical constituents and bioactive principles from the Mexican truffle and fermented products of the derived fungus *Ustilago maydis* MZ496986. *J. Agric. Food Chem.* 2023; 71(2): 1122-31.
- [53] Juárez-Montiel M, Ruiloba de León S, Chávez-Camarillo G, Hernández-Rodríguez C, Villa-Tanaca L. Huitlacoche (corn smut), caused by the phytopathogenic fungus *Ustilago maydis*, as a functional food. *Rev. Iberoam. Micol.* 2011; 28(2): 69-73.
- [54] Bautista-González J. A. Conocimiento tradicional de los hongos medicinales en seis localidades diferentes del país [Tesis de licenciatura]. México: Facultad de Ciencias. 2013.
- [55] Salazar López JM, Martínez Saldaña MC, Reynoso Camacho R, Chávez Morales RM, Sandoval Cardoso ML, Guevara Lara F. Antioxidant capacity and phytochemical characterization of ethanolic extracts from raw and cooked huitlacoche (*Ustilago maydis*-*Zea mays*) *Rev. Mex. Cienc. Farm.* 2017; 48(3): 37-47.
- [56] Bazzano LA, He J, Ogden LG, Loria CM, Vupputuri S, Myers L, et al. Fruit and vegetable intake and risk of cardiovascular disease in US adults: the first national health and nutrition examination survey epidemiologic follow-up Study. *Am. J. Clin. Nutr.* 2002; 76(1): 93-9.
- [57] Brighenti F, Valtueña S, Pellegrini N, Ardigò D, Del Rio D, Salvatore S, et al. Total antioxidant capacity of the diet is inversely and independently related to plasma concentration of high-sensitivity C-reactive protein in adult Italian subjects. *Br. J. Nutr.* 2005; 93(5): 619-25.
- [58] Pitsavos C, Panagiotakos DB, Tzima N, Chrysohoou C, Economou M, Zampelas A, Stefanadis C. Adherence to the Mediterranean diet is associated with total antioxidant capacity in healthy adults: the ATTICA study. *Am. J. Clin. Nutr.* 2005; 82(3): 694-99.
- [59] Katsnelson A, De Strooper B, Zoghbi HY. Neurodegeneration: from cellular concepts to clinical applications. *Sci. Transl. Med.* 2016; 8(364): 364ps18.
- [60] Martínez Leo EE, Segura Campos MR. Systemic oxidative stress: a key point in neurodegeneration- A review. *J. Nutr. Health Aging.* 2019; 23(8): 694-9.
- [61] Pegoretti V, Swanson KA, Bethea JR, Probert L, Eisel ULM, Fischer R. Inflammation and oxidative stress in multiple Sclerosis: consequences for therapy development. *Oxid. Med. Cell. Longev.* 2020; 12:(2020) 7191080.

- [62] Sienes Bailo P, Llorente Martín E, Calmarza P, Montolio Breva S, Bravo Gómez A, Pozo Giráldez A, et al. The role of oxidative stress in neurodegenerative diseases and potential antioxidant therapies. *Adv. Lab. Med.* 2022; 3(4): 351–60.
- [63] Dávila-Cervantes CA, Pardo-Montaña AM. Cambios en la esperanza de vida por causas de muertes crónicas en adultos mayores. México 2000-2013. *Rev. Cienc. Salud* 2017; 15(2): 223-35.
- [64] Roa Rojas PA, Martínez Ruiz A, Gutiérrez Robledo LM. Envejecimiento y demencias en México desde una perspectiva de género. *Género y salud* 2017; 15(3): 18.
- [65] Gómez Delgado Y, Velázquez Rodríguez EB. Health and food culture in Mexico. *RDU.* 2019; 20(1).
- [66] González-Cervantes ME, Hernández-Urbe JP, Gómez-Aldapa CA, Navarro-Cortez RO, Palma-Palma HM, Vargas-Torres A. Physicochemical, functional, and quality properties of fettuccine pasta added with huitlacoche mushroom (*Ustilago maydis*) *J. Food Process. Preserv.* 2021; 45(10): e15825.
- [67] Beas FR, Loarca PG, Guzmán MSH, Rodríguez MG, Vasco MNL, Guevara F. Potencial nutracéutico de componentes bioactivos presentes en huitlacoche de la zona centro de México. *Rev. Mex. Cienc. Farm.* 2011; 42(2): 36-44.
- [68] Estrada AF, Brefort T, Mengel C, Díaz-Sánchez V, Alder A, Al-Babili S, Avalos J. *Ustilago maydis* accumulates β -carotene at levels determined by a retinal-forming carotenoid oxygenase. *Fungal Genet. Biol.* 2009; 46(10): 803–13.
- [69] Vezza T, Canet F, De Maraón AM, Bañuls C, Rocha M, Víctor VM. Phytosterols: Nutritional health players in the management of obesity and its related disorders. *Antioxidants (Basel)* 2020; 9(12): 1266.
- [70] Ras RT, Trautwein EA. Consumer purchase behaviour of foods with added phytosterols in six European countries: Data from a post-launch monitoring survey. *Food Chem. Toxicol.* 2017; 110: 42–8.
- [71] Valadez Azúa R, Moreno Fuentes Á, Gómez Álvarez G. Cujtlacoche. El huitlacoche. 1ra ed. México: UNAM Instituto de investigaciones Antropológicas; 2011; 89: 104.
- [72] González A, Cruz M, Losoya C, Nobre C, Loreda A, Rodríguez R, Contreras J, Belmares R. Edible mushrooms as a novel protein source for functional foods. *Food Function* 2020; 23:11(9): 7400-14.
- [73] Scialabba NE-H. *Managing health livestock production and consumption.* Academic Press; 2022.
- [74] Santos A, Navascués E, Bravo E, Marquina D. *Ustilago maydis* killer toxin as a new tool for the biocontrol of the wine spoilage yeast *Brettanomyces bruxellensis*. *Int. J. Food Microbiol.* 2011; 31:145(1): 147-54.
- [75] Cortés-Sánchez A, Hernández-Sánchez H, Jaramillo-Flores M. Production of glycolipids with antimicrobial activity by *Ustilago maydis* FBD12 in submerged culture. *African J. Microbiol. Res.* 2011; 5: 2512–23.
- [76] Yang Long X, Takayoshi A, Wakimoto T, Abe I. Induced production of the novel glycolipid ustilagic acid C in the plant pathogen *Ustilago maydis*. *Tetrahedron Lett.* 2013; 54(28): 3655–57.
- [77] Becker F, Stehlik T, Linne U, Bölker M, Freitag J, Sandrock B. Engineering *Ustilago maydis* for production of tailor-made mannosylerythritol lipids. *Metab. Eng. Commun.* 2021; 12: e00165.
- [78] Steinberg G, Perez-Martin J. *Ustilago maydis*, A new model system of fungi for cell biology. *Trends Cell Biol.* 2008; 18(2): 61–7.