

This manuscript has been accepted after a double-blind peer review and will be published in a future issue of this journal. Its content is final and only the information regarding volume, number, publication date and pagination will change in the final version

## Complicated grief as a mental health sequel derived from the COVID-19 pandemic

### El duelo complicado como una secuela en salud mental derivada de la pandemia por COVID-19

Brenda Elizabeth Barraza-Sánchez<sup>a</sup>

---

#### Abstract:

The SARS-CoV2 pandemic is the third most serious coronavirus outbreak in less than 20 years, it has caused the death of millions of people around the world, mostly elderly people or those with a compromised immune status. This health emergency has become one of the main public health problems, precisely because of all the consequences to the physical and mental health of the population. For its part, complicated grief represents one of the most obvious consequences, due to the multiple losses experienced by the population and which could not be experienced normally due to health restrictions. However, until now, little research has been carried out in this regard. Until now, it has been determined that the participation of mental health professionals, psychologists and psychiatrists is essential for the diagnosis and care of complicated grief and other mental illnesses that may be a consequence of this pandemic, which is why this article has the objective to demonstrate the importance of the analysis and study of this phenomenon to develop effective interventions at the population and individual level.

#### Keywords:

Psychological attention, SARS-CoV2, complicated grief, mental health

---

#### Resumen:

La pandemia por SARS-CoV2 es el tercer brote más grave de coronavirus en menos de 20 años, ha provocado la muerte de millones de personas alrededor del mundo, en su mayoría personas de la tercera edad o con un estado inmunológico comprometido. Esta emergencia sanitaria se ha convertido en uno de los principales problemas de salud pública, precisamente por todas las consecuencias a la salud física y mental de la población. Por su parte, el duelo complicado representa una de las secuelas más evidentes, debido a las múltiples pérdidas que experimenta la población y que no pudieron ser vividas con normalidad debido a las restricciones sanitarias. Sin embargo, hasta este momento son pocas las investigaciones que se han realizado al respecto. Hasta ahora se ha determinado que es indispensable la participación de los profesionales de la salud mental, psicólogos y psiquiatras para el diagnóstico y atención de un duelo complicado y otros padecimientos mentales que pueden ser consecuencia de esta pandemia, por lo que este artículo tiene el objetivo de evidenciar la importancia del análisis y estudio de este fenómeno para desarrollar intervenciones eficaces a nivel poblacional e individual.

#### Palabras Clave:

Atención psicológica, SARS-CoV2, duelo complicado, salud mental

---

#### INTRODUCTION

The COVID-19 pandemic not only brought about damage to the physical health of humanity, but has also generated instability in public health and social systems due to the forced closure of schools, social institutions, as well as the collapse of the global economy due to the suspension of production, commercial and

labor activities.<sup>1</sup> Scientific research on epidemics and pandemics that have been carried out throughout human history has made it possible to understand that, during the outbreak of any infectious disease, the psychological reactions of the population play a fundamental role in the spread of a certain disease. pathology, in addition to the appearance of mental problems, such as anguish, fear, uncertainty, stress, depression


---

<sup>a</sup> Corresponding author, Consultorio privado | consultorio privado | Hidalgo | México, <https://orcid.org/0000-0001-1234-1234>, Email: barrabren18@gmail.com

and anxiety during and after the outbreak, and which are not exclusive to an age group or with certain cultural, economic or social characteristics, but it occurs in a generalized way.<sup>2</sup> Specifically speaking of the grief that people experience as a consequence of the COVID-19 pandemic, they not only refer to the death of a loved one, whether associated with SARS-CoV2 or any other cause, but humanity has been faced with various losses simultaneously due to restrictive and preventive measures to prevent the spread of the virus.<sup>1</sup> For this reason, it is necessary to consider various psychological factors that intervene in times of pandemic, firstly, trust in the management by the authorities and the health sector, of preventive measures that were implemented to avoid the risks of contracting the virus, as well as those aimed at the treatment and hospitalization of infected patients, ensuring dignified treatment, and on the other hand, those that correspond to community support in the different processes that were experienced such as confinement, changes in routines and lifestyles, and in those who have contracted the virus, facing isolation and in serious cases, hospitalization, solidarity and emotional displays adapted to health measures influence the person's ability to assimilate losses and cope with grief in a better way. All of these aspects have resulted in mental health conditions with the development of complicated grief, depressive and anxiety disorders, suicidal thoughts, sleep and appetite disorders.<sup>3</sup> Considering the relevance of the analysis of the mental health situation of the population and the consequences left by the pandemic, the objective of this article is to carry out a bibliographic review, which allows identifying the importance of the analysis and study of mental conditions such as complicated grief and that this information allows the development of effective interventions at the population and individual level.

### PSYCHOLOGICAL IMPACT OF A PANDEMIC

The history of humanity has been marked by a series of epidemics and pandemics over time, in figure 1. Main pandemics worldwide show some of the different periods in which diseases occurred that became epidemics and pandemics, claiming the lives of thousands and millions of people in various parts of the world.<sup>4</sup>



**Figure 1. Timeline, main epidemics worldwide.<sup>4</sup>**

The analysis through scientific research of these phenomena throughout history has made it possible to realize that, during any outbreak of an infectious disease, the psychological reactions of the population play a fundamental role in the spread of a certain pathology, in addition to the appearance of mental problems, such as emotional distress, fear, uncertainty, stress, depression and anxiety to name a few, during and after the health emergency and that are not exclusive to an age group or with certain cultural, economic or social characteristics, but it occurs in a generalized way.<sup>2</sup>

Many public health studies have shown that outbreaks of infectious diseases such as severe acute respiratory syndrome (SARS) in 2003<sup>5</sup>, the influenza A (H1N1) epidemic in 2009<sup>6</sup> and the Ebola outbreak in 2014, lead the population to experience severe psychological problems, such as post-traumatic stress disorder, depressive disorders and anxiety disorders, specifically generalized anxiety and social anxiety were identified.<sup>7,8</sup>

However, despite the high rates of mental health problems that arise in populations during epidemics or pandemics, few economic, material and human resources are allocated specifically for mental health care, inside and outside of the country hospitals, which may be understandable in these cases, since the priority is to reduce the spread of the disease, however, psychological and psychiatric needs should not be overlooked during any phase of pandemic management, since as mentioned, psychological factors play a fundamental role in compliance with health measures issued by governments and health organizations to avoid mass infections and likewise in how people face the threat of infection and the consequent losses.<sup>9</sup>

### TYPES OF GRIEF

The word mourning comes from the Latin *dolium*, which means pain or affliction. Also defined as the natural reaction to the loss of an object, event or person that is very significant, as well as those emotional reactions and behaviors derived from the breaking of an emotional bond. Therefore, grief considers physical, psychological and social factors in addition to their duration and intensity, since these are indicators of the dimension of the loss and are considered to evaluate whether it is a normal grieving process or a complicated one.<sup>10</sup>

According to the classifications of psychiatric medical manuals, grief can be normal and complicated, as explained in Table 1, which describes the characteristics that allow identifying each of them.

According to the Diagnostic and Statistical Manual of Mental Disorders (DSM-V) 11, normal grief is accompanied by some of the characteristic symptoms of a major depressive episode; however, they may be normal and understandable given the situation of significant loss, so an evaluation is necessary, especially considering the time of the loss in relation to these symptoms, to determine the presence of a mood disorder.


**Table 1.** Main characteristics of normal and complicated grief.<sup>10-12</sup>

	COMPLICATED GRIEF	NORMAL GRIEF
<b>Main manifestations</b>	Symptoms persist and become more severe over time. Other mental pathologies can develop, depressive disorders and anxiety disorders.	Intense and persistent feeling of sadness. Hopelessness. Anger and frustration. Isolation. Insomnia, loss of appetite and weight.
<b>Duration</b>	The symptoms last for more than 18 months and become more intense for the sufferer.	Approximately 6 to 12 months, although residual symptoms may occur after that time.
<b>Prevalence</b>	It occurs in approximately 10 to 20% of the general population.	At least once in their lives, everyone will experience a grieving process.

On the other hand, complicated grief implies that this prolongation of symptoms can affect going through the different stages of this process, accompanied by a lack of emotional regulation, which can generate, as a consequence, a series of maladaptive behaviors that lead to clinical symptoms. severe such as depression, anxiety and in severe cases psychotic outbreaks, these consequences will be given by elements such as the type of bond with the person who has been lost, the characteristics of the loss and the mental aspects of the bereaved, as well as social support.<sup>13, 14</sup>

**STAGES OF GRIEF**

Grief is a process that consists of a series of phases or stages, although this theory has not been unified, since its definition is conditioned according to clinical observation or the therapeutic approach, so there are several concepts that cannot be definitive.<sup>15</sup> The American psychiatrist Elisabeth Kubler-Ross and the medical Kessler<sup>16</sup> is one of the most prominent researchers on death and palliative care, she proposes five stages of grief. Each of these stages is mentioned below (figure 2).



**Figure 2.** Stages of grief.<sup>16</sup>

**PSYCHOLOGICAL INTERVENTION IN COMPLICATED GRIEF**

The first-line treatment for grief that becomes complicated or pathological is usually psychological therapy. Cognitive

behavioral therapy in this case promotes the person's adaptive change in the face of loss, by focusing on negative cognitions and behaviors<sup>14</sup>. Other investigations conclude in their research that cognitive behavioral therapy for the treatment of pathological grief has been more effective when carried out in a group.<sup>17</sup>

It was also found that telemedicine turned out to be a good option to care for those who were grieving during the pandemic, a series of small randomized controlled trials showed that cognitive behavioral therapy through electronic media and the internet was effective at that time for grief care and that people who required it could more easily access a psychiatric evaluation and diagnosis.<sup>18</sup>

On the other hand, there is evidence that psychotherapy during the grieving process that becomes complicated offers better results to reduce symptoms when it is the mourner himself who requests care and in the same way when it is identified that it is a complicated or complicated grief. that in addition to grief, some mental disorder such as depression, anxiety or post-traumatic stress occurs.<sup>19</sup>

Research indicates that group therapy can be a good option for the treatment of grief symptoms and that it can even help patients address grief that they experienced in other stages of their lives, allowing for a more adequate elaboration, which can mean in some cases the bases for their recovery.<sup>15</sup>

Regarding the way in which a psychotherapist could make a diagnosis of complicated grief, one can consider the Complicated Grief Inventory (CDI) adapted to Spanish, which is a self-administered questionnaire with 19 items with 5 Likert-type response categories that assess the frequency of the symptom explored in the emotional, cognitive or behavioral field. If 25 points or more are obtained, it is considered a complicated duel.<sup>20</sup>

Speaking specifically about the interventions that were carried out during the pandemic for grief care, the Ministry of Health<sup>21</sup> issued a document called recommendations for grieving family members during the COVID-19 pandemic, the objective of which was to issue guidelines for mourning. self-care in different contexts to prevent the spread of the virus, Table 2 describes the recommendations for the population in more detail.

**Table 2.** Self-care guidelines for grief during the COVID-19 pandemic.<sup>21</sup>

Care guidelines	Recommendations
Self-care	Sleep hygiene. Adequate nutrition. Avoid consumption of psychoactive substances.
Emotional	Use of electronic devices to communicate and express emotions. Respect moments of solitude, but without reaching isolation. If you require support for physical or emotional care, it is suggested to call the lines of authorized public institutions.
Cognitive	Do not demand too much of yourself intellectually, avoid the infodemic. Write objectives in relation to grief, in order to identify its emotional and physical evolution. Not making important decisions.
Spiritual	Create a special space where significant objects are placed to remember the person. Activities that enhance creativity. Farewell rituals from home, where they can be connected on electronic devices to be shared.

On the other hand, Spain issued a manual of recommendations for relatives of patients who died from complications derived from the virus, which, in the face of health isolation measures, in the most critical period in terms of infections and deaths, suggesting an adaptation of rituals and ceremonies inside the home and emphasized requesting support from mental health professionals through telephone lines specifically set up for this work in order to prevent complications in grief and likewise suicidal ideation or attempts.<sup>22</sup>

#### GRIEF IN TIMES OF COVID-19

Speaking specifically about losses and consequent grief during the COVID-19 pandemic, mourners were exposed to two main risk factors that could cause complicated grief, the first of which is related to those situational factors around the death, since that this, in most cases, arrives suddenly, also accompanied by feelings of helplessness, not being able to accompany the loved one during the illness process and without being able to say goodbye before death.<sup>23</sup>

The second refers to the circumstantial factors in which the loss occurs, in this case, the limitations that arose regarding social support during the grieving process, since this made it difficult for the mourner to express his feelings and share them with others. others to validate their pain, as well as remember memories and perform funeral rituals or ceremonies, all of this allows the mourners to assume the pain of their loss in solitude.<sup>24</sup> Regarding the relational factors, of the mourner with the deceased, and which in most cases are those that represent a greater risk, for example, the type of bond they maintained, if there was any type of dependence, economic, physical or emotionally, complicated grief is more likely to develop when a parent or partner is lost.<sup>17</sup>

Likewise, it is necessary to consider some individual aspects that could be considered to evaluate the presence of complicated grief, including previous unresolved grief, presence of any mental disorder prior to the loss, especially mental state disorders. mood, abuse of psychoactive substances, few emotional and social resources to cope with the loss.<sup>25</sup>

#### CONCLUSIONS

The COVID-19 pandemic is one of the worst global public health crises in a century, whose medical complications have claimed the lives of several million people around the world and left even more people grieving as a result. Although the possible medium-term consequences are just beginning to be identified and it is still unknown what will come in the long term after this health crisis, it has been determined that a significant number of grieving people will possibly develop complicated grief, derived from the health measures they implemented. in the face of emergency.<sup>2</sup>

Research has determined that if psychotherapeutic care is not received for complicated grief, the possibility of developing some mental pathology increases, including depression, anxiety, sleep disorders, eating disorders and a greater risk of suicide, in addition to increases the risk of developing cardiovascular and oncological diseases, as well as deficiencies in the immune system, complicated grief can also cause difficulties in adherence to treatment in patients who require daily medication, for example, those who suffer from chronic diseases.<sup>27</sup>

Likewise, it is essential that health professionals be able to detect symptoms of depression, especially if there is suicidal ideation and even suicide attempts derived from a significant loss, as well as a possible post-traumatic stress disorder in these patients, and consider referral to specialized psychiatric and psychological services in a timely manner.<sup>18</sup>

## REFERENCES

- [1] Ayittey FK, Ayittey MK, Chiwero NB, Kamasah JS, Dzuvor, C. Economic impacts of Wuhan 2019-nCoV on China and the world. *J. Med. Virol.* 2020; 92(1):473-5.
- [2] Cullen W, Gulati G, Kelly BD. Mental health in the COVID-19 pandemic. *QJM.* 2020; 113(5):311-2.
- [3] Organización Panamericana de la salud, Organización Mundial de la Salud. Intervenciones recomendadas en salud mental y apoyo psicosocial durante la pandemia [SMAPS]. [Internet]. [location unknown]: 2020. [cited 2024 february 24] Available from: [https://iris.paho.org/bitstream/handle/10665.2/52427/OPSNMHNMCOVID-19200026\\_spa.pdf?sequence=1&isAllowed=y](https://iris.paho.org/bitstream/handle/10665.2/52427/OPSNMHNMCOVID-19200026_spa.pdf?sequence=1&isAllowed=y)
- [4] Talevi D, Socci V, Carai M, Carnaghi G, Faleri S, Trebbi E, et al. Mental health outcomes of the CoViD-19 pandemic. *Riv. Psichiatri.* 2020; 55(3):137-44.
- [5] Peng EYC, Lee MB, Tsai ST, Yang CC, Morisky DE, Tsai LT. et al. Population-based post-crisis psychological distress: An example from the SARS outbreak in Taiwan. *J. Formos Med. Assoc.* 2010; 109(7):524-32.
- [6] Taha SA, Matheson A, Anisman H. H1N1 was not all that scary: uncertainty and stressor appraisals predict anxiety related to a coming viral threat. *Stress Health* 2014; 30(2):149-57.
- [7] Ehlers A, Clark DM. A. Cognitive model of post-traumatic stress disorder. *Behav. Res. Ther.* 2000; 38(4):319-45.
- [8] Shultz JM, Baingana F, Neria Y. The Ebola Outbreak and Mental Health in 2014: Current Status and Recommended Response. *JAMA*, 2015; 313(6):567-8.
- [9] Taylor S. *The Psychology of Pandemics: Preparing for the Next Global Outbreak of Infectious Disease.* UK: Cambridge Scholars Publishing; 2019.
- [10] Meza EG, García S, Torres A, Castillo L, Suárez SS, Martínez S. El proceso del duelo. Un mecanismo humano para el manejo de las pérdidas emocionales. *Rev. Esp. Méd. Quir.* 2008; 13(1):28-31.
- [11] American Psychiatric Association. *Guía de consulta de los criterios diagnósticos del DSM-5®: Spanish Edition of the Desk Reference to the Diagnostic Criteria From DSM-5®.* American Psychiatric Pub; 2014.
- [12] Hamui P. El duelo en tiempos de coronavirus (COVID-19). *Gravida.* [Internet]. [location unknown]: 2020. [cited 2024 february 24] Available from: <https://spm.mx/el-duelo-en-tiempos-de-coronavirus-covid-19/>
- [13] Millan-González R, Solano-Medina N. Duelo, duelo patológico y terapia interpersonal. *Rev. Colombia Psiqu.* 2010; 39(2):375-88.
- [14] Eisma MC, Boelen PA, Lenferink LI. Prolonged grief disorder following the coronavirus (COVID-19) pandemic. *Psychiatry.* Res. 2020; 288:113031.
- [15] Gamó E, Pazos P. El duelo y las etapas de la vida. *Rev. Asoc. Esp. Neuropsi.* 2009; 29(104):455-69.
- [16] Kübler-Ross E, Kessler D. *On grief and grieving: finding the meaning of grief through the five stages of loss.* New York: Scribner; 2007.
- [17] Iglewicz A, Shear MK, Reynolds CF, Simon N, Lebowitz B. Complicated grief therapy for clinicians: an evidence-based protocol for mental health practice. *Depress. Anxiety* 2020; 37(1): 90-8.
- [18] Yahya AS, Khawaja S. Bereavement and Grief During the COVID-19 Pandemic. *Prim. Care Companion CNS Disord.* 2020; 22(4):20com02661.
- [19] Allumbaugh DL, Hoyt WT. Effectiveness of grief therapy: A meta-analysis. *J. Couns. Psych.* 1999; 46(3):370-80.
- [20] Limonero JT, Lacasta M, García, JA, Maté, J, Prigerson, HG. Adaptación al castellano del inventario de duelo complicado. *Med. Paliativ.* 2009; 16(5):291-7.
- [21] Gobierno de México. Recomendaciones para familiares en duelo durante la pandemia de COVID-19. [Internet]. [location unknown]: 2020. [cited 2024 february 24] Available from: [https://coronavirus.gob.mx/wpcontent/uploads/2020/06/SaludMental\\_FamiliaresEnDuelo.pdf](https://coronavirus.gob.mx/wpcontent/uploads/2020/06/SaludMental_FamiliaresEnDuelo.pdf)
- [22] Sociedad Española de Cuidados Paliativos (SECPAL). Recomendaciones para familiares para la despedida y el duelo ante la presencia del COVID-19. [Internet]. [location unknown]: 2020. [cited 2024 february 23] Available from: <http://www.secpal.com/Documentos/Blog/covid19%20duelo%20y%20despedia%20familiares%20documento%20final%20V2.pdf>
- [23] Lobb E, Kristjanson L, Aoun S, Monterosso L, Halkett G, Davies A. Predictors of complicated grief: a systematic review of empirical studies. *Death Stud.* 2010; 34(8):673-98.
- [24] Larrotta-Castillo R, Méndez-Ferreira AF, Mora-Jaimes C, Córdoba-Castañeda MC, Duque-Moreno J. Pérdida, duelo y salud mental en tiempos de pandemia. *Rev. Univ. Indust. Santander* 2020; 52(2):179-80.
- [25] Kokou-Kpolou CK, Fernández-Alcántara M, Cénat JM. Prolonged Grief Related to COVID-19 Deaths: Do We Have to Fear a Steep Rise in Traumatic and Disenfranchised Grievs? *Psychol. Trauma.* 2020; 12(S1):S94-S95.
- [26] Gesi C, Carmassi C, Cerveri G, Carpita B, Cremona IM, Dell'Osso L. Complicated Grief: What to Expect After the Coronavirus Pandemic. *Front. Psychiatry* 2020; 11:489.
- [27] Osiris RM, Guerrero GE, Herrera A, Ascencio L, Rodríguez ME, Aguilar I, et al. *Manual operativo para el curso de acompañamiento del duelo en situaciones especiales durante la pandemia de la COVID-19 en México.* 2020.