

Current trends and future directions of mHealth in psychology: Challenges and promising perspectives

Tendencias actuales y futuras de mHealth en psicología: Desafíos y perspectivas prometedoras

Luis A. Rivera-Ramírez^a

Abstract:

The aim of the following text is to explore the importance of mobile health (mHealth) applications in mental health, addressing their integration into traditional clinical settings, their effectiveness in diverse populations, and the ethical challenges associated with their use. It seeks to understand how mHealth interventions can complement and reinforce the work of mental health specialists, as well as identify the specific needs of different groups. In addition, it aims to highlight the importance of considering ethical aspects such as privacy, data security and fairness in access to these technologies. It seeks to provide a comprehensive view of the benefits, challenges, and future directions in the use of mobile health applications to improve psychological care.

Keywords:

e-health, m-health, intervention, mental health, psychology

Resumen:

El objetivo del siguiente texto es explorar la importancia de las aplicaciones de salud móvil (mHealth) en la salud mental, abordando su integración en entornos clínicos tradicionales, su efectividad en diversas poblaciones y los desafíos éticos asociados con su uso. Se busca comprender cómo las intervenciones de mHealth pueden complementar y reforzar el trabajo de los especialistas en salud mental, así como identificar las necesidades específicas de diferentes grupos. Además, se pretende destacar la importancia de considerar aspectos éticos como la privacidad, la seguridad de los datos y la equidad en el acceso a estas tecnologías. Se busca proporcionar una visión integral de los beneficios, desafíos y futuras direcciones en el uso de aplicaciones de salud móvil para mejorar la atención psicológica.

Palabras Clave:

e-health, m-health, intervención, salud mental, psicología

INTRODUCTION

E-health interventions, like m-health or mobile health, have gained a lot of attention for their potential to address psychological problems. The World Health Organization defines e-health as the use of Information and Communication Technologies (ICTs) in support of health and health-related areas.¹ This term encompasses a wide range of applications, including telemedicine, interventions, human and economic capital management, just to mention a few.² On the other hand, mobile health can be defined as the use of mobile technologies (cell phones, tablets, wearables, among others) for health purposes.¹

The benefits of e-health and mHealth interventions include their availability, accessibility, cost-effectiveness, scalability, and ability to personalize content.³ In addition, these interventions provide real-time strategies and can calibrate the intensity of the intervention according to the needs of the users.⁴ They also have the potential to expand the scope of psychology services, improving the health and well-being of children, young people and their families, as well as positively intervening on sleep quality, fatigue and physical activity in patients and survivors of some diseases such as cancer.⁴⁻⁶

Additionally, the use of mHealth interventions has been associated with reductions in stress and depressive symptoms, higher levels of emotional well-being, and improved mental health outcomes in diverse populations.^{6,7} That is why e-health

^a Corresponding author, Universidad Autónoma del Estado de Hidalgo | Subentidad | Pachuca-Hidalgo | México, <https://orcid.org/0000-0003-2095-2765>, Email: luis_rivera@uaeh.edu.mx

and mHealth interventions have shown promise in supporting health specialists.

IMPACT OF MHEALTH APPS ON MENTAL HEALTH

The use of mobile applications has a significant impact on health, as they complement and reinforce the work of the specialist. However, it is important to note that they do not completely replace the assistance and accompaniment of a health professional.⁸

The literature highlights the importance of integrating psychological theories into mHealth applications to improve their impact and adherence to mental health interventions.⁹ The most commonly used psychological approach in this type of intervention is cognitive-behavioral therapy, as it offers various techniques that can be self-administered (self-recording thoughts, beliefs, emotions, breathing techniques, among others) with appropriate training.¹⁰ In addition, it has been observed that these interventions can have long-term effects, such as the sustainability of healthy behaviors and the continuous improvement of mental health.^{11,12}

However, the need for more research to comprehensively assess the impact of mHealth apps on mental health, especially in specific populations such as cancer patients, older adults, and healthcare workers is recognized.¹³⁻¹⁵ Furthermore, it is emphasized the importance of considering sociocultural experiences and user preferences when developing mHealth applications for mental health.^{16,17}

ACCESSIBILITY AND REACH OF PSYCHOLOGICAL SERVICES THROUGH MHEALTH

The accessibility and scope of psychological services through mHealth is a very important point, due to the viability of the technology. In spite of acknowledging the potential benefits of mHealth in providing psychological interventions to diverse populations, such as children, adolescents or college students¹⁸, mHealth adoption remains a challenge in these environments.¹⁹ This is because, despite being constant users of the internet and mobile apps, they usually don't use health apps.²⁰

It is essential to understand the factors that influence the quality, regulation, and scope of virtual psychological counseling services.²⁰ Some studies have shown that knowledge about mHealth among university students in developing countries is low, and there are barriers to its use, such as ethnicity.²¹ For example, in countries such as Ghana, Benin, and Nigeria, where cell phone penetration is high, many people are unaware of the benefits these devices can provide to improve their health. In addition, the usability of mobile health apps varies, with some studies indicating low usability scores.²²

The potential of mHealth to support community health workers in delivering psychological interventions in underserved rural areas is highlighted.¹⁸ In addition, the use of mHealth has been identified as a viable modality to help vulnerable populations manage their chronic conditions, including mental health conditions.²³ However, the effectiveness of mHealth

interventions in developing countries, especially in the context of psychological services, remains unclear.²⁴

A relevant point to ensure the success of interventions through mHealth is to consider the commitment and motivation that the user has to adhere to the solutions offered by those who develop the intervention programs.²⁵ It is also necessary to take into account the design principles (which are mentioned in the section on user-centric application development and design) of mHealth interventions to achieve sustainable changes in health behaviors.²⁶

The development and evaluation of mHealth apps for psychological care, especially for vulnerable populations such as pregnant women, people with disabilities, ethnicity communities, migrants, and others, require specialized frameworks to ensure the quality and effectiveness of the apps.²⁷ Therefore, while mHealth holds promise for expanding the accessibility and reach of psychological services, especially in developing countries, there are challenges related to adoption, usability, and efficacy that need to be addressed. Understanding the specific needs of target populations and developing personalized mHealth solutions, along with rigorous assessment and regulation, are crucial to harnessing the full potential of mHealth in the provision of psychological interventions.

ETHICAL CONSIDERATIONS IN THE USE OF MHEALTH

The use of mHealth in psychology raises important ethical considerations, especially with regard to privacy, safety, and quality of interventions. The privacy policies of mHealth apps are essential, as they collect and store personal data, including sensitive information. This requires clear and comprehensive privacy policies to protect user data.²⁸ For example, policies that ensure data encryption, limit access to sensitive information, and establish clear protocols for handling sensitive data.

Mobile health research, both regulated and unregulated, presents ethical challenges that require measures to protect end-users and ensure the benefits of it.²⁹ Ethical frameworks for planning, implementation, and evaluation are essential, especially in sensitive areas such as treatment adherence. Considerations such as participant autonomy, informed consent, and personal data protection must be integrated throughout the process.³⁰

Ensuring privacy and security in mHealth applications requires critical criteria and countermeasures to safeguard user data and maintain trust in the use of these technologies.³¹ Understanding the barriers and facilitators of mobile health applications from a security perspective is crucial to address ethical concerns and ensure the safe and effective use of these technologies.³² For example, implementing robust security protocols such as two-factor authentication and limited access to sensitive data can help protect users' privacy and mitigate security risks. Furthermore, it is imperative to set the agenda for bioethics in the mobile healthcare revolution to address ethical challenges and protect vulnerable groups from harm and discrimination.³³ It is crucial to consider the circumstances of vulnerable groups, such as those

with limited access to resources or facing systematic discrimination, in order to implement measures that protect their integrity and rights.

These references collectively underscore the ethical complexities surrounding the use of mHealth in psychology, emphasizing the need for robust privacy policies, regulatory frameworks, and security measures to ensure the ethical use of mobile technologies in psychological interventions. Identifying and addressing potential risks from a bioethical standpoint is essential to ensure equity and protection for all participants.

USER-CENTRIC APPLICATION DEVELOPMENT AND DESIGN

The design of mental health apps exerts a significant influence on user experience and adherence to interventions.¹ First, it highlights the importance of developing the application based on user-centered design principles, which refer to an approach that puts the end user in the design process, ensuring that the interface and functionalities are tailored to the needs and preference of the target users.³⁴

Usability, which refers to the ease of use and effectiveness of the app, while accessibility which refers to the app's ability to be used by people with different skills and needs, are crucial factors that contribute to user engagement and adherence to interventions.³⁵ To do this, it is necessary to ensure intuitive navigation and a clear presentation of information. Incorporating principles for digital program development, aligned with best practices in user-centered design, is considered essential to creating user-friendly mental health apps.³⁵

The app should prioritize maintaining the user-provider relationship, fostering a sense of connection and support, especially in the context of mental health care.³⁴ Furthermore, the development of interventions that promote user engagement and strengthen care networks is vital for enhancing the user experience.

INTEGRATION OF MHEALTH IN TRADITIONAL CLINICAL SETTINGS

The clinical setting is defined as the space where tasks, activities, and processes aimed at the care, promotion, protection, and restoration of the health of a defined population are carried out.³⁶ It is within this context that mHealth has been documented in various healthcare settings, spanning from disease diagnosis to medication referral, medical education, and disease management for patients.³⁷ These apps seek to address the unmet needs of health systems by providing tools that empower healthcare providers, disseminate health information, and facilitate contact tracing during a health crisis such as pandemics.³⁸

MHealth has been shown to support the performance of healthcare workers by providing clinical updates, learning materials, and reminders, especially in underserved rural areas.¹⁸ It is important to note that conventional treatment interventions using technology alone may have limited efficacy. For example, it was observed that an intervention carried out entirely with

technology, applied in the context of pediatric obesity, faced problems of adherence to treatment³⁹

However, mHealth applications have demonstrated efficacy in promoting changes in health behaviors and improving health outcomes by providing quick access to health information and facilitating interactive interventions.⁴⁰ Furthermore, technology-based remote interventions, such as the use of phones, pedometers, and internet-based methods, have shown promise in promoting physical activity compared to minimal treatments.⁴¹ Healthcare provider training has also been extended through the use of mHealth applications, utilizing mobile phones as innovative work assistants for community healthcare professionals.⁴² The prioritization of mobile health in healthcare models has been observed in countries with higher life expectancies and the need for continuous patient monitoring.⁴³ Notwithstanding these advancements, the impact of mobile medical photography applications on clinical care is still unclear, highlighting the need for further research in this area.⁴⁴ As mHealth continues to expand, it enables healthcare providers to deliver better care to patients in remote and underserved communities, thus addressing the challenges of healthcare accessibility.⁴⁵ Mobile and web applications offer opportunities for promoting oral health, indicating the potential of mobile health to contribute to preventive healthcare measures.⁴⁶

It is crucial to recognize the substantial benefits of conventional practices and manual therapy in healthcare, especially in the context of stroke rehabilitation.⁴⁷ The potential of mHealth applications to monitor overweight and obesity within the Internet of Things framework has been proposed, highlighting the role of technology in promoting healthy behaviors and providing automated interventions.⁴⁸

The effectiveness of mobile health technologies in enhancing healthcare services has been highlighted, with emerging literature on the application of mobile technologies in low-income countries.⁴⁹ It is crucial for mHealth applications to incorporate evidence-based approaches to minimize the risk of harm and ensure the safe application of knowledge by users.⁵⁰ Personalized healthcare systems facilitated by mHealth services and applications enhance communication between patients and healthcare providers, thereby contributing to improved health outcomes.⁵¹

MHEALTH & SPECIFIC POPULATIONS

The technology has been widely used in the field of psychology, aimed at addressing the specific needs of diverse populations, including adolescents, older adults, women, and individuals with particular disorders. A systematic review on digital mental health interventions for depression, anxiety, and improving psychological well-being among college students. Most of the selected studies were conducted in the United States, United Kingdom, Ireland, Australia, Canada and China with a total of 89 studies included in the analysis. We found that 47% of the interventions evaluated showed effectiveness, while 34% demonstrated partial effectiveness. These results underscore the

significant potential of these types of interventions to address the mental health needs of young university students globally.⁵²

In a gender-responsive study conducted in Brazil, a wellness app designed to provide relaxation training and positive psychology principles was evaluated, with the goal of reducing chronic stress and improving the overall well-being of working women. After the implementation of the intervention, the experimental group had a 17% increase in subjective well-being (measured by the WHO-5 scale), compared to the control group.⁵³

In another study carried out in China, a mobile app intervention was developed for pregnant women, with the aim of training them in mindfulness to mitigate perinatal depression and other mental health problems. The study involved 1,140 women. As a result, it was observed that the group that received the intervention experienced a 60.9% reduction in the risk of depressive symptoms compared to those that did not receive the training through the mobile app.⁵⁴

Mobile-enhanced single-session interventions in severe mental illness have been investigated using a randomized controlled trial, indicating the potential of brief mobile interventions to address mental health challenges. The impact of cell phone support on the psychosocial outcomes of young people living with HIV and non-adherence to antiretroviral therapy has also been studied, demonstrating the potential of mobile interventions to support people with specific health conditions.⁵⁵

APPS CREATED IN MEXICO

Despite the advances in the use of mobile applications to promote people's well-being, and the presence of a wide variety of these in the various application stores, there is scarce scientific evidence base related to mobile applications designed and applied specifically in the Mexican population. Table 1 summarizes the interventions carried out in Mexico.

Next, the first app titled 'What happens if you go too far?'⁵⁶, This program aimed to prevent substance use in adolescents and was implemented on a high school campus as part of the introductory curriculum for the school year. A pretest-posttest design was applied and a Google form was used to assess life skills. Participants received training on the effects and risks of psychoactive substance use, along with strengthening life skills. The intervention was complemented with strategies at home, and had a total duration of 10.5 hours distributed in group sessions.

This intervention sought to equip adolescents with skills to resist peer pressure and make healthy choices regarding substance use. It is highlighted that the use of mobile technologies in intervention processes can expand the scope of preventive work among adolescents. The next intervention, called RatinARBox⁵⁷, It aimed to develop an augmented reality environment to teach operant conditioning to psychology students. In this research, group video calls were held to download the application and evaluations were made regarding the experience. A significant improvement was observed in students who used the app compared to those who did not. Participants highlighted the usefulness and interactivity of the application. It is concluded that

the RatinARBox application is a promising tool to strengthen the understanding of operant conditioning concepts in psychology students, demonstrating its potential in technology-based education.

Table 1. mHealth interventions carried out in the Mexican population.⁵⁸⁻⁶⁰

App Name	Population	Target
"What happens if you go too far?"	Adolescents (12-18 years)	Increase the perception of risk of tobacco, alcohol and marijuana use
RatInARBox	Students (average age 18)	Augmented Reality Environment for Teaching Operant Conditioning
Live Tobacco-Free... Make up Your Mind!	Adults (>18 years)	Smoking Cessation Plan

The third intervention or application carried out in Mexico corresponds to an intervention carried out on adults with the name 'Live without Tobacco...'⁵⁸, with the purpose of helping participants quit smoking. Although it was a pilot study, it was widely accepted by participants, generating high levels of satisfaction and high participation throughout the 12-week program. At the end of the program, it was verified through biochemical examinations that 40% of the participants had quit smoking. These findings suggest that mobile interventions may be an effective tool to promote smoking cessation in underserved communities, highlighting the importance of accessibility and personalization in health care.

These intervention examples demonstrate that mobile apps have a lot to offer in the health arena, from addressing addictive behaviors to providing training for students. We are just at the beginning of an advance in quality and acceptance by the Mexican population in relation to these applications."

CHALLENGES AND FUTURE DIRECTIONS

Mobile technology in health has emerged as a promising field in psychology, providing innovative interventions to address challenges in mental health service delivery. However, there are current challenges and areas that require further research in this field, as well as possible future directions and advancements.

These include measuring user engagement in eHealth and mHealth behavior change interventions.⁵⁹ In addition, it points to the need for strategies for research in mHealth, as researchers may face unique challenges in this line of research. It also highlights the importance of understanding the information

needs of informal caregivers in the management of behavioral and psychological symptoms of people with dementia, as well as the design of mHealth applications to address these needs.⁶⁰ Another crucial challenge is the inclusion of people with disabilities in mHealth, as its omission could lead to health disparities.⁶¹

It is essential to recognize the existence of the digital divide, which affects marginalized social, cultural, and demographic groups, leading to inequalities in access, trust, and technological skills.⁶² Although modern life is closely tied to cell phone use, we cannot ignore the reality of those who face difficulties accessing technology. To achieve equitable and effective implementation, it is crucial to address and reduce the impact of technology exclusion. In this sense, digital literacy, which implies the ability to use, understand and leverage ICTs effectively, emerges as an essential element to close the digital divide and promote technological inclusion.⁶³

CONCLUSION

In today's era, technology has become an indispensable part of our lives, offering unprecedented opportunities for healthcare professionals. mHealth emerges as a crucial tool in this landscape, providing additional support and facilitating access to healthcare for patients. In the field of psychology, although there has been some resistance towards the adoption of these interventions due to ignorance and perceived risks, it is important to recognize their potential to provide relief to a greater number of people and offer complementary alternatives in patient health care. It is essential to be open to exploring and taking advantage of the opportunities that technology offers us in the continuous improvement of health care and the well-being of our patients.

REFERENCES

- [1] World Health Organization [Internet]. Directriz de la OMS: Recomendaciones sobre intervenciones digitales para fortalecer los sistemas de salud, Geneva: World Health Organization. 2022 - [cited 2024 Feb 12]. Available from: <https://iris.who.int/bitstream/handle/10665/354399/9789240046993-spa.pdf?sequence=1>.
- [2] Díaz de León Castañeda C, Góngora Ortega J. eSalud en servicios de salud públicos en México: estudio de caso. *Reg. Soc.* 2020;32:e1256.
- [3] Velandia Bernal VB, Lozano Rodríguez M, Baquero Mujica GK [Internet]. Aplicaciones móviles en salud, una revisión sistemática cualitativa, Universidad de La Salle. 2021 - [cited 2024 Feb 12]. Available from: <https://ciencia.lasalle.edu.co/cgi/viewcontent.cgi?article=2910&context=optometria>.
- [4] Clough BA, Casey LM. The smart therapist: A look to the future of smartphones and mHealth technologies in psychotherapy. *Prof. Psychol. Res. Pr.* 2015;46(3):147-53.
- [5] Jukic T, Ihan A, Strojnik V, Stubljarić D, Starc A. The effect of active occupational stress management on psychosocial and physiological well-being: a pilot study. *BMC. Med. Inform. Decis. Mak.* 2020;20(321):1-8.
- [6] Kosasih FR, Yee VTS, Toh SHY, Sündermann O. Efficacy of Intellect's self-guided anxiety and worry mobile health programme: A randomized controlled trial with an active control and a 2-week follow-up. *PLOS Digit. Health* 2023;2(5):e0000095.
- [7] Oliveira C, Pereira A, Vagos P, Nóbrega C, Gonçalves J, Afonso B. Effectiveness of Mobile App-Based Psychological Interventions for College Students: A Systematic Review of the Literature. *Front. Psychol.* 2021;12:(647606):1-16.
- [8] Izquierdo Alonso JL, Almonacid Sánchez C. Nuevas tecnologías en Medicina. *RIECS* 2022;7(1):69-82.
- [9] Denecke K, Schmid N, Nüssli S. Implementation of Cognitive Behavioral Therapy in e-Mental Health Apps: Literature Review. *J. Med. Internet Res.* 2022;24(3):e27791.
- [10] Rathbone AL, Clarry L, Prescott J. Assessing the Efficacy of Mobile Health Apps Using the Basic Principles of Cognitive Behavioral Therapy: Systematic Review. *J. Med. Internet Res.* 2017;19(11):e399.
- [11] Vázquez-de Sebastián J, Ciudin A, Castellano-Tejedor C. Analysis of Effectiveness and Psychological Techniques Implemented in mHealth Solutions for Middle-Aged and Elderly Adults with Type 2 Diabetes: A Narrative Review of the Literature. *J. Clin. Med.* 2021;10(12):1-23.
- [12] Wong AKC, Wong FKY, Bayuo J, Chow KKS, Wong SM, Lee AYL. A randomized controlled trial of an mHealth application with nursing interaction to promote quality of life among community-dwelling older adults. *Front. Psychiatry* 2022;13:1-12.
- [13] Akingbade O, Adediran V, Somoye IE, Alade AS, Chow KM. Perceived feasibility and usefulness of mHealth interventions for psychoeducational support among Nigerian women receiving chemotherapy for breast cancer: a focus group study. *Support. Care Cancer* 2022;30(12):9723-34.
- [14] Akinsulore A, Aloba O, Oginni O, Oloniniyi I, Ibigbami O, Seun-Fadipe CT, et al. Developing an mHealth intervention to reduce COVID-19-associated psychological distress among health care workers in Nigeria: Protocol for a design and feasibility study. *JMIR Res. Protoc.* 2022;11(11):e36174.
- [15] Walsh JC, Richmond J, Mc Sharry J, Groarke A, Glynn L, Kelly MG, et al. Examining the impact of an mHealth behavior change intervention with a brief in-person component for cancer survivors with overweight or obesity: Randomized controlled trial. *JMIR MHealth UHealth* 2021;9(7):e24915.
- [16] Willis HA, Neblett EW. Developing culturally-adapted mobile mental health interventions: a mixed methods approach. *Mhealth* 2023;9(1):1-25.
- [17] Müssener U, Thomas K, Bendtsen P, Bendtsen M. High school students' preferences and design recommendations for a mobile phone-based intervention to improve psychological well-being: Mixed methods study. *JMIR Pediatr. Parent.* 2020;3(2):e17044.
- [18] Källander K, Tibenderana JK, Akpogheneta OJ, Strachan DL, Hill Z, ten Asbroek AHA, et al. Mobile health (mHealth) approaches and lessons for increased performance and retention of community health workers in low- and middle-income countries: A review. *J. Med. Internet Res.* 2013;15(1):e17.
- [19] Hoque MR. An empirical study of mHealth adoption in a developing country: the moderating effect of gender concern. *BMC. Med. Inform. Decis. Mak.* 2016;16(1):1-10.
- [20] Torres Escobar GA, Giraldo Roncancio L, Restrepo Benavides PA, Giraldo Benavides AK, Giraldo Benavides LD, Montenegro Cedeño DC, et al. Reflexiones sobre el bienestar universitario virtual en Colombia. *Incl. Desarro.* 2019;6(2):23-33.

- [21] Peprah P, Abalo EM, Agyemang-Duah W, Gyasi RM, Reforce O, Nyonyo J, et al. Knowledge, attitude, and use of mHealth technology among students in Ghana: A university-based survey. *BMC Med. Inform. Decis. Mak.* 2019;19(1):1-11.
- [22] Arafa A, Mostafa ZM, Sheerah HA, Alzahrani F, Almuzaini Y, Senosy S, et al. MHealth app barriers, usability, and personalization: A cross-sectional study from Egypt and Saudi Arabia. *J. Pers. Med.* 2022;12(12):1-11.
- [23] Humble JR, Tolley EA, Krukowski RA, Womack CR, Motley TS, Bailey JE. Use of and interest in mobile health for diabetes self-care in vulnerable populations. *J. Telemed Telecare* 2016;22(1):32-8.
- [24] Mao Y, Lin W, Wen J, Chen G. Impact and efficacy of mobile health intervention in the management of diabetes and hypertension: a systematic review and meta-analysis. *BMJ Open Diabetes Res. Care* 2020;8(1):e001225.
- [25] Mustafa AS, Ali N, Dhillon JS, Alkaws G, Baashar Y. User engagement and abandonment of mHealth: A cross-sectional survey. *Healthcare* 2022;10(2):1-13.
- [26] Yang L, Kuang A, Xu C, Shewchuk B, Singh S, Quan H, et al. Design principles in mHealth interventions for sustainable health behavior changes: Protocol for a systematic review. *JMIR Res. Protoc.* 2023;12:e39093.
- [27] Ayyoubzadeh SM, Ahmadi M, Khounraz F, Ahmadi M, Pourhamidi R, Abbasi S. Requirement analysis of mHealth app for pregnancy care and a framework suggestion for evaluating pregnancy mobile apps. *J. Obstet. Gynecol. Cancer Res.* 2022;7(6):554-62.
- [28] Sunyaev A, Dehling T, Taylor PL, Mandl KD. Availability and quality of mobile health app privacy policies. *J. Am. Med. Inform. Assoc.* 2015;22(e1):e28-33.
- [29] Hammack-Aviran CM, Brelsford KM, Beskow LM. Ethical considerations in the conduct of unregulated mHealth research: Expert perspectives. *J. Law Med. Ethics* 2020;48(1 suppl):9-36.
- [30] DiStefano MJ, Schmidt H. mHealth for Tuberculosis Treatment Adherence: A Framework to Guide Ethical Planning, Implementation, and Evaluation. *Glob. Health Sci. Pract.* 2016;4(2):211-21.
- [31] Rezaee R, Khashayar M, Saedinezhad S, Nasiri M, Zare S. Critical criteria and countermeasures for mobile health developers to ensure mobile health privacy and security: Mixed methods study. *JMIR MHealth UHealth* 2023;11:e39055.
- [32] Zhou L, Bao J, Watzlaf V, Parmanto B. Barriers to and facilitators of the use of mobile health apps from a security perspective: Mixed-methods study. *JMIR MHealth UHealth* 2019;7(4):e11223.
- [33] Lucivero F, Jongsma KR. A mobile revolution for healthcare? Setting the agenda for bioethics. *J. Med. Ethics* 2018;44(10):685-9.
- [34] Silva-Vázquez PO, Benítez-Guerrero E, Rosales-Morales VY. Hacia un modelo para la generación automática de interfaces gráficas centradas en el usuario. *Atas da conferência Ibero-Americana IADIS* 2019;325-28.
- [35] Prefasi Gomar S, Magal-Royo T. Principios de Diseño en las aplicaciones de m-Salud Mental orientadas al control psicosocial. *INNODOCT* 2020;413-20.
- [36] Ayala G, Jiménez R, Jiménez C. El proceso de integración del alumno al entorno clínico para el aprendizaje reflexivo. *ENEO-UNAM* 2010;7(4):53-61.
- [37] Mosa ASM, Yoo I, Sheets L. A systematic review of healthcare applications for smartphones. *BMC Med. Inform. Decis. Mak.* 2012;12(1):1-31.
- [38] Davalbhakta S, Advani S, Kumar S, Agarwal V, Bhoyar S, Fedirko E, et al. A systematic review of smartphone applications available for corona virus disease 2019 (COVID19) and the assessment of their quality using the Mobile Application Rating Scale (MARS). *J. Med. Syst.* 2020;44(9):1-15.
- [39] Kouvari M, Karipidou M, Tsiampalis T, Mamalaki E, Poulimeneas D, Bathrellou E, et al. Digital health interventions for weight management in children and adolescents: Systematic review and meta-analysis. *J. Med. Internet Res.* 2022;24(2):e30675.
- [40] Hakala S, Rintala A, Immonen J, Karvanen J, Heinonen A, Sjögren T. Effectiveness of physical activity promoting technology-based distance interventions compared to usual care. Systematic review, meta-analysis and meta-regression. *Eur. J. Phys. Rehabil. Med.* 2017;53(6):953-67.
- [41] Rickard N, Arjmand HA, Bakker D, Seabrook E. Development of a Mobile Phone App to Support Self-Monitoring of Emotional Well-Being: A Mental Health Digital Innovation. *JMIR Ment. Health.* 2016;3(4):e49.
- [42] Shawky M, Salem MR, Abouhashima F, Abdelaziz S, Aguizy FHE. Use of a mHealth approach for the training of health-care providers on nutrition counseling in a malnutrition clinic. *Open Access Maced. J. Med. Sci.* 2022;10(E):1350-8.
- [43] Moreira R, Teles A, Fialho R, Baluz R, Santos TC, Goulart-Filho R, et al. Mobile applications for assessing human posture: A systematic literature review. *Electronics* 2020;9(8):1-24.
- [44] Wyatt KD, Willaert BN, Lohse CM, Pallagi PJ, Yiannias JA, Hellmich TR. Experiences of health care providers using a mobile medical photography application. *Appl. Clin. Inform.* 2020;11(1):122-9.
- [45] Wright V, Dalwai M, Smith RV, Jemmy JP. Médecins Sans Frontières' Clinical Guidance mobile application: analysis of a new electronic health tool. *Public Health Action* 2015;5(4):205-8.
- [46] Rasmus K, Toratti A, Karki S, Pesonen P, Laitala ML, Anttonen V. Acceptability of a mobile application in children's oral health promotion-A pilot study. *Int. J. Environ. Res. Public Health* 2021;18(6):1-9.
- [47] Selamat SNS, Che Me R, Ahmad Ainuddin H, Salim MSF, Ramli HR, Romli MH. The Application of Technological Intervention for Stroke Rehabilitation in Southeast Asia: A Scoping Review with Stakeholders' Consultation. *Front. Public Health* 2022;9:1-12.
- [48] Alturki R, Alharbi AI, AlQahtani SS, Khan S, Truong Hoang V. A proposed application for controlling overweight and obesity within the framework of the Internet of things. *Mob. Inf. Syst.* 2021; 2021:1-9.
- [49] Free C, Phillips G, Felix L, Galli L, Patel V, Edwards P. The effectiveness of M-health technologies for improving health and health services: a systematic review protocol. *BMC Res. Notes* 2010;3:1-7.
- [50] Gladman T, Tylee G, Gallagher S, Mair J, Rennie SC, Grainger R. A tool for rating the value of health education mobile apps to enhance student learning (MARuL): Development and usability study. *JMIR MHealth UHealth* 2020;8(7):e18015.
- [51] Marques G, Pitarma R. MHealth: Indoor environmental quality measuring system for enhanced health and well-being based on internet of things. *J. Sens. Actuator Netw.* 2019;8(3):1-20.
- [52] Lattie EG, Adkins EC, Winquist N, Stiles-Shields C, Wafford QE, Graham AK. Digital mental health interventions for depression, anxiety,

and enhancement of psychological well-being among college students: Systematic review. *J. Med. Internet Res.* 2019;21(7):e12869.

- [53] Coelho CC, Toba PR, Lacerda SS, Lima AH, Barrichello CRC, Amaro Jr E, et al. A new mental health mobile app for well-being and stress reduction in working women: Randomized controlled trial. *J. Med. Internet Res.* 2019;21(11):e14269.
- [54] Sun Y, Li Y, Wang J, Chen Q, Bazzano AN, Cao F. Effectiveness of smartphone-based mindfulness training on maternal perinatal depression: Randomized controlled trial. *J. Med. Internet Res.* 2021;23(1):e23410.
- [55] Sayegh CS, MacDonell KK, Clark LF, Dowshen NL, Naar S, Olson-Kennedy J, et al. The impact of cell phone support on psychosocial outcomes for youth living with HIV nonadherent to antiretroviral therapy. *AIDS Behav.* 2018;22(10):3357-62.
- [56] Fuentes A PMDC, Jiménez Tapia A, Ruiz-Cortés EM, Bolaños-Ceballos F, Flores Castro JC, Gutiérrez R, González-Forteza C. Effectiveness of a Mobile App to Increase Risk Perception of Tobacco, Alcohol, and Marijuana Use in Mexican High School Students: Quantitative Study. *JMIR Mhealth Uhealth* 2023;11:e37873.
- [57] Sánchez-Sordo JM, Teodoro-Vite S. Desarrollo de un entorno de realidad aumentada para la enseñanza del condicionamiento operante en Psicología. *Tecnol. Cienc. Educ.* 2022;23:115-36.
- [58] Cupertino AP, Cartujano-Barrera F, Ramírez M, Rodríguez-Bolaños R, Thrasher JF, Pérez-Rubio G, Falfán-Valencia R, Ellerbeck EF, Reynales-Shigematsu LM. A Mobile Smoking Cessation Intervention for Mexico (Vive sin Tabaco... ¡Decídete!): Single-Arm Pilot Study. *JMIR Mhealth Uhealth* 2019;7(4):e12482.
- [59] Short CE, DeSmet A, Woods C, Williams SL, Maher C, Middelweerd A, et al. Measuring engagement in eHealth and mHealth behavior change interventions: Viewpoint of methodologies. *J. Med. Internet Res.* 2018;20(11):e292.
- [60] Jagoda T, Dharmaratne S, Rathnayake S. Informal carers' information needs in managing behavioural and psychological symptoms of people with dementia and related mHealth applications: a systematic integrative review to inform the design of an mHealth application. *BMJ Open* 2023;13(5):e069378.
- [61] Jones M, Morris J, Deruyter F. Mobile Healthcare and People with Disabilities: Current State and Future Needs. *Int. J. Environ. Res. Public Health.* 2018;15(3):1-13.
- [62] Hardy A, Ward T, Emsley R, Greenwood K, Freeman D, Fowler D, et al. Bridging the digital divide in psychological therapies: Observational study of engagement with the SlowMo mobile app for paranoia in psychosis. *JMIR Hum. Factors* 2022;9(3):e29725.
- [63] Alarcón Belmonte I, Sánchez Collado R, Yuguero O, Acezat Oliva J, Martínez-Millana A, Saperas Pérez C. La alfabetización digital como elemento clave en la transformación digital de las organizaciones en salud [Digital literacy as a key element in the digital transformation of health organizations]. *Aten. Primaria* 2024;56(6):1-6.