

**Canadian Foundation for Dietetic Research
Dietetic Research Event – June 14-15, 2012**

These abstracts represent research projects of dietitians that were accepted through a peer-review process for presentation.

*Indicates the presenter [R] = Research abstract [E] = Experience-sharing abstract

**e-POSTER RESEARCH PRESENTATION ABSTRACTS
THURSDAY JUNE 14 , 2012**

CLINICAL RESEARCH, INCLUDING OUTCOMES OF INTERVENTION

Body mass index, waist circumference, risk eating behaviors and attitudes toward body figure in a female sample of Moncton University students.

T J Saucedo-Molina*1, L Villalon *2, J Zaragoza Cortes *1, A Peña-Irecta *1.

1Autonomous University of Hidalgo. Academic Area of Nutrition. Hidalgo, Mexico.

2Université de Moncton. École de sciences des aliments, de nutrition et d'études familiales. Faculté des sciences de la santé et des services communautaires. Moncton, New Brunswick. [R]

Introduction: The multifactorial origin of eating disorders is well known and among the most studied risk factors are: eating behaviors, high body mass index (BMI) and body thin-ideal internalization. Several studies have confirmed a high prevalence of these risk factors, particularly in adolescents and young adults, and predominantly in females. Objective: The purpose of this study was to evaluate the prevalence of risk eating behaviors and body thin-ideal internalization, and its relationship with BMI and 11 waist circumference (WC) in a sample of female students from Moncton University in Canada. Method: A sample of 84 female students from Moncton University with a range of 18 to 30 years ($X=21\pm 2.1$), completed self-reported multidimensional questionnaire to assess risk eating behaviors and body thinideal internalization. To calculate Body Mass Index (BMI), each subject was weighed and measured, as well to get WC. Results: Results showed that 75 % had a normal weight. There were fewer students in the very underweight (1%) and underweight (3%) categories, than those in the overweight (15%) and obese (6%) categories. The prevalence of risk eating behaviors was 6% (mainly dietary restraint) meanwhile body thin-ideal internalization was 29%. For both risk factors, the analysis by BMI showed that the higher percentages were obtained for the overweight students. From 14% of students who reached $WC>80$ cm, 50% reported risk eating behaviors, and 70% body thin-ideal internalization. Significant correlations were founded between risk eating behaviors, attitudes toward body figure, BMI and WC. Conclusions: These findings are a significant contribution in the research field of eating disorders in Moncton, Canada and it must be consider in the design of new research in this topic.